

Statin Drugs May Lead To Co-Q10 Deficiency, Diabetes Risk

Main Category: [Statins](#)

Also Included In: [Diabetes](#)

Article Date: 14 Apr 2013 - 0:00 PDT

A laboratory study has shown for the first time that coenzyme Q10 offsets cellular changes that may be linked to a side-effect of some statin drugs - an increased risk of adult-onset [diabetes](#).

Current ratings for:

Statin Drugs May Lead To Co-Q10 Deficiency, Diabetes Risk

Patient / Public:
Healthcare Prof:



4.2 (5 votes)
4.75 (4 votes)

Statins are some of the most widely prescribed drugs in the world, able to reduce LDL, or "bad" [cholesterol](#) levels, and the risk of [heart attacks](#) or other cardiovascular events. However, their role in raising the risk of diabetes has only been observed and studied in recent years.

The possibility of thousands of statin-induced diabetics is a growing concern, and led last year to new labeling and warnings by the Food and Drug Administration about the drugs, especially when taken at higher dosage levels.

The findings of the new research were published as a rapid communication in *Metabolic Syndrome and Related Disorders*, and offer another clue to a possible causative mechanism of this problem.

Pharmacy researchers at Oregon State University who authored the study said the findings were made only in laboratory analysis of cells, and more work needs to be done with animal and ultimately human studies before recommending the use of coenzyme Q10 to help address this concern.

"A number of large, randomized clinical trials have now shown that use of statins can increase the risk of developing type-2 diabetes by about 9 percent," said Matthew K. Ito, an OSU professor of pharmacy and president-elect of the National Lipid Association.

"This is fairly serious, especially if you are in the large group of patients who have not yet had a cardiovascular event, but just take statin drugs to lower your risks of [heart disease](#)," Ito said.

A suspect in this issue has been altered levels of a protein called GLUT4, which is part of the cellular response mechanism, along with insulin, that helps to control blood sugar levels. A reduced expression of GLUT4 contributes to insulin resistance and the onset of type-2 diabetes, and can be caused by the use of some statin drugs.

The statins that reduce cholesterol production also reduce levels of coenzyme Q10, research has shown. Coenzyme Q10 is needed in cells to help create energy and perform other important functions. And this study showed in laboratory analysis that if coenzyme Q10 is supplemented to cells, it prevents the reduction in GLUT4 induced by the statins.

Not all statin drugs, however, appear to cause a reduction in GLUT4.

The problems were found with one statin, simvastatin, that is "lipophilic," which means it can more easily move through the cell membrane. Some of the most commonly used statins are lipophilic, including simvastatin, [atorvastatin](#), and lovastatin. All of these statins are now available as generic drugs, and high dosage levels have been most often linked with the increase in diabetes.

Tests in the new study done with a "hydrophilic" statin, in this case pravastatin, did not cause reduced levels of GLUT4. Pravastatin is also available as a generic drug.

"The concern about increasing levels of diabetes is important," Ito said. "We need to better understand why this is happening. There's no doubt that statins can reduce cardiovascular events, from 25-45 percent, and are very valuable drugs in the battle against heart disease. It would be significant if it turns out that use of coenzyme Q10 can help offset the concerns about statin use and diabetes."

Before that conclusion can be reached, the researchers said, additional studies are needed on coenzyme Q10 supplementation and the pathogenesis of statin-induced diabetes.

-
-
-

Article adapted by Medical News Today from original press release. Click 'references' tab above for source. Visit our [statins](#) section for the latest news on this subject.

Add Your Opinion On This Article

'Statin Drugs May Lead To Co-Q10 Deficiency, Diabetes Risk'

Please note that **we publish your name**, but we **do not publish** your email address. It is only used to let you know when your message is published. We do not use it for any other purpose. Please see our [privacy policy](#) for more information.

If you write about specific medications or operations, please **do not** name health care professionals by name.

All opinions are moderated before being included (to stop spam)

Contact Our News Editors

For any corrections of factual information, or to contact the editors please use our [feedback form](#).

Please send any medical news or health news press releases to: pressrelease@medicalnewstoday.com

Note: Any medical information published on this website is not intended as a substitute for informed medical advice and you should not take any action before consulting with a health care professional. For more information, please read our [terms and conditions](#).